Environmental Protection Agency

(d) Indirect or inadvertent residues. [Reserved]

[45 FR 55198, Aug. 19, 1980, as amended at 63 FR 10543, Mar. 4, 1998; 63 FR 65073, Nov. 25, 1998; 66 FR 28672, May 24, 2001; 68 FR 37764, June 25, 2003; 68 FR 48312, Aug. 13, 2003]

§ 180.396 Hexazinone; tolerances for residues.

(a) General. Tolerances are established for combined residues of the herbicide hexazinone (3-cyclohexyl-6-(dimethylamino)-1-methyl-1, 3, 5-triazine-2,4(1*H*,3*H*)-dione) and its metabolites (calculated as hexazinone) in or on the following food commodities:

| Commodity | Parts per million |
|-------------------------|----------------------|
| Alfalfa green forage | 2.0 |
| Alfalfa, hay | 8.0 |
| Blueberry | 0.2 |
| Cattle, fat | 0.1 |
| Cattle, meat byproducts | 0.1 |
| Cattle, meat | 0.1 |
| Goat, fat | 0.1 |
| Goat, meat byproducts | 0.1 |
| Goat, meat | 0.1 |
| Grass, pasture | 10 |
| Grass, range | 10 |
| Hog, fat | 0.1 |
| Hog, meat byproducts | 0.1 |
| Hog, meat | 0.1 |
| Horse, fat | 0.1 |
| Horse, meat byproducts | 0.1 |
| Horse, meat | 0.1 |
| Milk | 0.1 |
| Pineapple (whole fruit) | 0.5 |
| Sheep, fat | 0.1 |
| Sheep, meat byproducts | 0.1 |
| Sheep, meat | 0.1 |

(b) Section 18 emergency exemptions. [Reserved]

(c) Tolerances with regional registrations. A tolerance with regional registration, as defined in §180.1(n) and which excludes use of hexazinone on sugarcane in Florida, is established for combined residues of the herbicide hexazinone (3-cyclohexyl-6-(dimethyamino)-1-methyl-1,3,5-tri-azine-2,4(1*H*,3*H*)-dione) and its metabolites (calculated as hexazinone) in or on the following food commodities:

| Commodity | Parts per million |
|-----------------|----------------------|
| Sugarcane, cane | 0.2 5.0 |

(d) Indirect or inadvertent residues. [Reserved]

[65 FR 33713, May 24, 2000]

§ 180.399 Iprodione; tolerances for residues.

(a) General. (1) Tolerances are established for the combined residues of the fungicide iprodione [3-(3,5-dichlorophenyl)-N-(1-methylethyl)-2,4-dioxo-1-imidazolidinecarboxamide], its isomer 3-(1-methylethyl)-N-(3,5-dichlorophenyl)-2,4-dioxo-1-imidazolidinecarboxamide, and its metabolite 3-(3,5-dichlorophenyl)-2,4-dioxo-1-imidazolidine-carboxamide in or on the following food commodities:

| Commodity | Parts per million |
|-----------------------------|----------------------|
| Almond, hulls | 2.0 |
| Almond | 0.3 |
| Apricot | 20.0 |
| Bean, dried, vine hay | 90.0 |
| Bean, dry, seed | 2.0 |
| Bean, forage | 90.0 |
| Bean, succulent | 2.0 |
| Blueberry | 15.0 |
| Boysenberry | 15.0 |
| Broccoli | 25.0 |
| Caneberries | 25.0 |
| Carrot, roots | 5.0 |
| Cherry (sweet), postharvest | 20.0 |
| Cherry, tart | 20.0 |
| Cotton, undelinted seed | 0.10 |
| Currant | 15.0 |
| Garlic | 0.1 |
| Ginseng, dried root | 4.0 |
| Ginseng, root | 2.0 |
| Grape | 60.0 |
| Grape, raisin | 300 |
| Kiwifruit | 10.0 |
| Lettuce | 25.0 |
| Nectarine, postharvest | 20.0 |
| Onion, dry bulb | 0.5 |
| Peach, postharvest | 20.0 |
| Peanut | 0.5 |
| Peanut, hay | 150.0 |
| Peanut hay | 150.0 |
| Plum, postharvest | 20.0 |
| Plum, prune | 20.0 |
| Potato | 0.5 |
| Raspberry | 15.0 |
| Rice, bran | 30.0 |
| Rice, grain | 10.0 |
| Rice, hulls | 50.0 |
| Rice, straw | 20.0 |
| Strawberry | 15.0 |

(2) Tolerances are established for the combined residues of iprodione [3-(3,5-dichlorophenyl)-*N*-(1-methylethyl)-2,4-dioxo-1-imidazolidinecarboxamide], its isomer [3-(1-methylethyl)-*N*-(3,5-dichlorophenyl)-2,4-dioxo-1-imidazolidinecarboxamide, and its metabolites [3-(3,5-dichlorophenyl)-2,4-dioxo-1-imidazolidine-carboxamide] and [*N*-(3,5-dichloro-4-hydroxyphenyl)-ureido-carboxamide], all expressed as iprodione equivalents in or on the following food commodities of animal origin: